



ICHEP 2014

3-10 July 2014 Valencia, Spain
Japan timezone

Overview

Scientific Programme

Call for Abstracts

View my contribution

Submit a new contribution

Timetable

My conference

My contributions

Book of abstracts

Conference Home Page

[Support](#)

[PDF](#)

Development of liquid scintillator containing zirconium complex for neutrinoless double beta decay experiment

Submitted by **Yoshiyuki Fukuda** on **11 Apr 2014** at **10:19**

Id: 255

Last modification: 9 May 2014 19:53

Contribution type: Poster

Content

An organic liquid scintillator containing zirconium complex was studied for neutrinoless double beta decay (ZICOS) experiment. A ^{96}Zr nuclei has a large Q-value (3.35MeV), and no experiment is planned to use as a target. In order to realize ton scale target isotope with good energy resolution (4% @2.5MeV), we have used zirconium beta-diketon complex which has huge solubility (over 10w.t.%) to the Anisole. However, the absorption wavelength of diketon ligand overlaps with the luminescence from Anisole. Therefore, the light yield of liquid scintillator decreased in proportion to the concentration of beta-diketon complex. In order to avoid this problem, we synthesized beta-keto ester complex introducing -OC₃H₇ or -OC₂H₅ substituent groups in the beta-diketon complex. These complexes have shorter absorption wavelength (245nm) than the emission wavelength of Anisole (275nm). The scintillation light yield recovered about double, however, did not reach at the expected value, because the residual absorption around the 275nm still exists. We have found that those were caused by the impurities of beta-keto ester complex and it will be solved by the purifying the complex. We obtained that the light yield was about 40% with respect to the original cocktail, and the energy resolution was 13% @1MeV even though 8.5% of PMT photo coverage for 10 w.t.% concentration, which corresponds to 4.1% @2.5MeV assuming 40% photo coverage for ZICOS detector. We have also found that a diethyl malonate ligand shifted the absorption peak to around 210nm, and the complex should have no quenching for the Anisole based liquid scintillator. Here we will report the present status of ZICOS project.

Comments

This project is supported by by Grant-in-Aid for Scientific Research (C) (No. 24540295) and Grant-in-Aid for Scientific Research on Innovative Areas (No. 26105502) from the Japanese Society for the Promotion of Science (JSPS).

Abstract status

ACCEPTED as Poster

Primary authors

Yoshiyuki Fukuda (Miyagi University of Education)

Co-authors

Shigetaka Moriyama (Kamioka Observatory, ICRR, University of Tokyo)
Izumi Ogawa (Faculty of Engineering, Fukui University)

Presenters

Yoshiyuki Fukuda (Miyagi University of Education)